
[How Obesity Can Rewire the Immune System](#)

[jchen](#)

Submitted by jchen on March 30, 2022

```
var addthis_share = { templates: { twitter: "When mice with atopic dermatitis—a common type of allergic skin inflammation—are treated with drugs that target the immune system, their thickened, itchy skin generally heals quickly." } }
```

When mice with atopic dermatitis—a common type of allergic skin inflammation—are treated with drugs that target the immune system, their thickened, itchy skin generally heals quickly. **Sagar Bapat, MD, PhD** and **Alex Marson, MD, PhD**, Director of the Gladstone-UCSF Institute of Genomic Immunology and affiliate faculty at Diabetes Center, along with colleagues at Salk Institute, **Ye Zheng, PhD**, an Associate Professor, and **Ronald Evans, PhD**, Director of Gene Expression Laboratory and the March of Dimes Chair in Molecular and Developmental Biology, have now discovered that the same treatment in obese mice makes their skin worse instead. That is because obesity changes the molecular underpinnings of allergic inflammation, both in mice and humans.

The new study in [Nature](#) shed light on how obesity can change the immune system and, potentially, how clinicians might be able to better treat allergies and asthma in obese people.

Sagar Bapat speaks with research associate Sarah Gayer in his lab in Diabetes Center. At left is graduate student Ben Lesch. *Image by Noah Berger*

During his graduate studies at Salk and subsequent research in Marson Lab, Bapat wanted to know how obesity affected atopic dermatitis. He discovered that when mice were made obese by eating a high-fat diet prior to the induction of dermatitis, they developed more severe disease than lean animals. To understand why, he and his colleagues analyzed the immune cells and molecules that were active in each group of mice.

“What we were expecting to see in the obese mice was just a greater degree of the same kind of inflammation,” says Bapat. “Instead, we saw a completely different kind of inflammation.”

The body’s helper T cells, which help protect against infection but also become overactive in autoimmune disease or allergies, can be grouped into three classes: TH1, TH2, and TH17 cells. Scientists had considered atopic dermatitis a TH2 disease; that means the TH2 cells are the ones causing the skin inflammation.

In lean mice with atopic dermatitis, Bapat and his colleagues indeed found that the TH2 cells were active. In obese mice with the same condition, however, TH17 cells were activated. At a molecular level, it meant that atopic dermatitis was completely different in the obese mice, raising the question of whether the drugs that work in lean animals would also be effective in obese animals.

A more extensive article from which this was adapted may be viewed [here](#).

[Research](#)

[News](#)